

[epsng,psng,jmm
 [1] 1 [1] / [1] 1 textbf{cmbxti10} textbf{ss} cmssbx10 mathbf{cmbxti10} mathbf{ss} cmssbx10 upmath
 eurm10 AMSa msam10 0upmath19 0upmath16 0upmath40 3AMSa36 3AMSa3E
 normalcmrmit boldcmrbxit [1] 1 [1] 1 [1] 1 normalcmrbxit boldcmrbxit normalcmssbxn bold-
 cmssbxn @mathgroup@group @mathgroup@normal@groupeurmn @mathgroup@bold@groupeurn @math-
 group@group @mathgroup@normal@groupmsamn @mathgroup@bold@groupmsamn @group19 @group16
 @group40 @group36 @group3E
 [1] 1 [1] 1 [1] 1 OT1cmrbxit boldOT1cmrbxit OT1cmssbxn boldOT1cmssbxn UPMUeurmn UPM-
 boldUeurbn AMSaUmsamn $\Gamma 0UPM$ $19\Gamma 0UPM$ $16\Gamma 0UPM$ $40\Lambda 3AMSa$ $36\Lambda 3AMSa$ $3E$

A New approach for a Galactic Synchrotron Polarized Emission Template in the Microwave Range [G. Bernardi et al.] G. Bernardi,^{1,2} E. Carretti,¹ R. Fabbri,³ C. Sbarra,¹ S. Poppi,⁴ S. Cortiglioni¹

24 October 2002

firstpage–lastpage 0000

document

firstpage

abstract We present a new approach in modelling the polarized Galactic synchrotron emission in the microwave range (20-100 GHz), where this radiation is expected to play the leading role in contaminating the Cosmic Microwave Background (CMB) data. Our method is based on real surveys and aims at providing the real spatial distributions of both polarized intensity and polarization angles. Its main features are the modelling of a polarization horizon to determine the polarized intensity and the use of starlight optical data to model the polarization angle pattern. Our results are consistent with several existing data, and our template is virtually free from Faraday rotation effects as required at frequencies in the cosmological window.